

# Scientific References

**1-** Probiotics to improve testicular function (Andrology 5:439–444, 2017). K. Tremellen and 2 K. Pearce.

<https://onlinelibrary.wiley.com/doi/pdf/10.1111/andr.12400>

**2-** Testosterone regulates keratin 33B expression in rat penis growth through androgen receptor signaling. Yan-Min Ma, Kai-Jie Wu. Qiang Dang.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4236322/>

**3-** Boas M, Boisen KA, Virtanen HE, Kaleva M, Suomi AM, et al. Postnatal penile length and growth rate correlate to serum testosterone levels: a longitudinal study of 1962 normal boys. Eur J Endocrinol. 2006;154:125–9.

<https://pubmed.ncbi.nlm.nih.gov/16382001/>

**4-** Camurdan AD, Oz MO, Ilhan MN, Camurdan OM, Sahin F, et al. Current stretched penile length: cross-sectional study of 1040 healthy Turkish children aged 0 to 5 years. Urology. 2007;70:572–5.

<https://pubmed.ncbi.nlm.nih.gov/17905119/>

**5-** Husmann DA. Micropenis: an animal model and its human correlates. Adv Exp Med Biol. 2002;511:41–54.

<https://pubmed.ncbi.nlm.nih.gov/12575755/>

**6-** Bin-Abbas B, Conte FA, Grumbach MM, Kaplan SL. Congenital hypogonadotropic hypogonadism and micropenis: effect of testosterone treatment on adult penile size why sex reversal is not indicated. J Pediatr. 1999;134:579–83.

<https://pubmed.ncbi.nlm.nih.gov/10228293/>

**7-** Tietjen DN, Uramoto GY, Tindall DJ, Husmann DA. Micropenis in hypogonadotropic hypogonadism: response of the penile androgen receptor to testosterone treatment. J Urol. 1998;160:1054–7.

<https://pubmed.ncbi.nlm.nih.gov/9719275/>

**8 -**Zhang YL, Zhou ZX, Zhang YD, Parker MG. Expression of androgen receptors and prostatic steroid-binding protein during development of the rat ventral prostate. J Endocrinol. 1988;117:361–6.

<https://pubmed.ncbi.nlm.nih.gov/3392492/>

**9-** Chang C, Saltzman A, Yeh S, Young W, Keller E, et al. Androgen receptor: an overview. Crit Rev Eukaryot Gene Expr. 1995;5:97–125.

<https://pubmed.ncbi.nlm.nih.gov/8845584/>